Key to <i>Juniperus</i> of the Far East (Japan, Taiwan, Sakhalin Island) rev. 2/2011 1a. Leaves all acicular (subulate, joined the base), in whorls of 3, spreading, articulate at base;
flowers axillary
2a. Leaves with 1 white stomatal band on the adaxial surface
3a. Leaves 4 - 10 (-12) mm x 1 mm or more wide, nearly flat to curved in cross section,
white stomatal band wider than green margins
3b. Leaves 7 - 25 mm x 0.5 - 0.9 mm wide, 'U' or 'V' shaped in cross section, with a keel
(ridge) on the abaxial side, deeply grooved, white stomatal band narrower that green
margins
4a. Upright shrub or tree; dry areas in mountains; cones 6-9 mm; leaves 12-25 mm. long
J. rigida var. rigida
4b.Low procumbent shrub on sands at seashores; cones 8-12 mm; leaves 10-22 mm. long
2b. Leaves with 2 white stomatal bands on the adaxial surface
5a. Leaves with mucronate tips, flat to U shaped, 12 - 20 mm; Taiwan, e. China
J. formosana
5b. Leaves with acute to obtuse, boat shaped, 7 - 12 mm, Japan
6a. Leaves with acute to obtuse tips, boat-shaped, 8 - 12 (-14) mm; endemic to Bonin
Islands
6b. Leaves acute to apiculate tips; boat-shaped, 7 - 10 mm and wide (1.1 - 1.5 mm),
endemic to Ryukyu Islands and s. coast of Shizuoka Pref., Oshima (Island), Japan
J. taxifolia var. lutchuensis
1b. Leaves not acicular (jointed at the base), but decurrent (with a sheath and blade), scale-like
and whip (decurrent with tips free), flowers terminal
7a All leaves whip type (decurrent with tips free), no scale-like leaves
8a. Procumbent shrubs, (1-) 2 - 3 seeds per cone, sea shores, Japan and Korea
J. procumbens
8b. Erect shrub or tree, 1 seed per cone; rocky cliffs, endemic to Taiwan J. morrisonicola
7b. Both scale-like and whip (decurrent with free tips) leaves present; whip-leaves
interspersed among sections of scale-leaves in mature branchlets, (1-) 2 seeds per cone
If almost all leaves scale-like, then a decumbent shrub (var. taiwanensis and var.
tsukusiensis)